





## Skin antisepsis

## Spray disinfection



Perform hygienic hand disinfection and put on low-germ examination gloves.



Immediately before operating, trim the hair using clean, sharp clippers. Avoid wet shaving to prevent skin injuries.

Use a paper towel or a (contactless) vacuum to remove the trimmed hair.



Clean dirt and oil from skin and decontaminate it before applying the antiseptic. Do this by rubbing cleansing foam into the skin and cleaning it.

Goal = removing larger impurities and transient skin flora.



After one minute, any foam residue can be removed with a clean cloth.
Bring the patient into the OR. Position the patient safely for the procedure.



Before applying the antiseptic, disinfect hands and put on sterile surgical gloves.



Aseptic: Goal 1) rapid, significant reduction of resident skin flora; 2) long-lasting (remanent) aseptic effect.

Spray Braunoderm® on the skin until it is thoroughly saturated.

## Skin antisepsis

## Spray disinfection



Use a sterile swab to wipe off any remaining Braunoderm®, moving from the center to the periphery. Use a new sterile swab each time. Repeat this process a total of three times.



Finish by moistening the skin once more with Braunoderm®.



After the exposure time has elapsed, the surgeon may apply a sterile dressing to the patient.



Additional information, this poster, and much more can be found at: www.bbraun-vetcare.com/inyourhands

B. Braun Vet Care GmbH | Am Aesculap-Platz | 78532 Tuttlingen | info@bbraun-vetcare.de | www.bbraun-vetcare.com

Thank you very much - The pictures were taken in Hergolding with the generous support of the small animal specialists Frontier GmbH.